

Using Apatite (U-Th)/He Thermochronometry to Study the Low-Temperature Thermal History of Shell Canyon, Northwestern Bighorn Mountains, Wyoming

Grant Kaye

Geology Department, The Colorado College 14. E. Cache La Poudre St., Colorado Springs, CO, 80903
Faculty Sponsors: Chris Siddoway, The Colorado College, Peter Crowley, Amherst College, Peter Reiners, Washington State University

INTRODUCTION

The Bighorn Mountains of north-central Wyoming provide an excellent opportunity to study the uplift and exhumation of crystalline basement in the Rocky Mountain region. (U-Th)/He dating, a recently refined (Wolf et al., 1996; House et al. 1997)

technique for studying the low temperature thermal history of apatite-bearing rocks, provides important constraints on the timing and geometry of uplift and exhumation of crust. The structure of the Bighorns and their marginal basins typifies mountain ranges uplifted in the Rocky Mountain region during the Laramide Orogeny, which lasted from ~75 to ~45Ma (Snook, 1992). Crustal shortening during this period caused Precambrian crystalline rocks to bulge upwards beneath Paleozoic and younger sediments, causing the sedimentary layers to dip steeply along the margins of the uplifted crystalline rocks into what are now the Powder River and Bighorn basins. The general structure of the Bighorns resembles a large-scale asymmetric anticline (see Figure 3). Easily accessible outcrops of Precambrian crystalline rock are found today throughout the range along US Route 14 through Shell Canyon over Granite Pass. Many petrologic and structural studies have been done on the Bighorn Mountains (e.g., Darton, 1906; Wise and Obi, 1992). Despite the extensive research done on the Bighorns and other Laramide ranges, little is known about their low-temperature exhumation history, and the (U-Th)/He ages obtained in this study provide valuable details about the timing and geometry of uplift and exhumation.




Figure One. Shaded 3D view up Shell Canyon to the east, showing samples, roads, and major streams. Scale varies in this perspective.

METHODS

Field – Bighorns

Samples collected from the Bighorn Mountains come from five geographic areas (Reuter, Fig. 2, this volume). They are (Table 1): an east-side elevation transect, a west-side elevation transect through Shell Canyon, a central plateau group consisting of range-parallel equal elevation samples, a group collected near Five Springs (northwest part of the range), and samples from the gneissic rocks comprising the southern part of the range. Because accurate location and elevation information are crucial in using He dates to constrain orogenic events, both GPS and an aneroid altimeter were used to locate the samples (~2-3 kg each) and their elevations. The location of each sample with respect to the Precambrian/Cambrian unconformity was also noted when possible. “BH##” samples were processed by 1999 Bighorns Keck Project participants, while “BHM-###” samples were He dated from separates provided generously by Dr. Robert Giegengack of the University of Pennsylvania.

Laboratory – Caltech

At Caltech, (U-Th)/He apatite separates were prepared from samples using standard techniques. Samples were crushed, pulverized, washed, and sieved to 50 and 230 mesh. Apatite was separated using standard magnetic and heavy liquid techniques. Only euhedral, inclusion-free grains (~80 microns in width and ~200 microns in length), were selected for dating. Typically, six to seven grains per sample were selected, measured, weighed, and placed in stainless steel vessels for ^4He analysis. ^4He contents were determined by outgassing the samples at 950°C, spiking with ^3He , and measuring $^4\text{He}/^3\text{He}$ on a quadrupole mass spectrometer in the Noble Gas Isotope lab of Kenneth Farley. Samples were then dissolved in HNO_3 , and U and Th concentrations were determined by isotope dilution on a sector ICP-MS.

Inclusions (such as zircon) present problems for U-Th/He analyses (Wolf et. al., 1996) by introducing anomalously high amounts of apparently parentless ^4He , giving inaccurately old ages. Extreme care in selecting inclusion-free grains was taken by scrutinizing them under cross-polarized light for uniform and invariant extinction. Also, during ^4He analysis, the “re-extract” process served to eliminate dates calculated from samples whose apatite grains possibly contained previously undetected inclusions. Apatite should completely outgas all ^4He during heating to ~950°C for about 20 minutes, but in some instances, additional ^4He was released from grains during a second heating step. This “re-extracted” helium may have originated from an inclusion of a mineral other than apatite in the sample grains. Ages obtained from samples that experienced “re-extracts” during analyses were discarded. When possible, additional grains were chosen from the remaining separates of samples, and additional He dating was performed on the second aliquot of apatite grains.

An additional correction was applied in the age calculation of each sample to account for loss of energetic ^4He from small grains (F_1 correction, Farley et al., 1996). ^4He can be lost disproportionately from alpha decay of ^{238}U atoms situated at the outer margins of an apatite crystal. F_1 values for each sample, indicative of the degree of the correction necessitated by the size of the sample grains, are given in column four of Table 1.

Region	Sample	He Age (Ma)	Elev. (m)	Age Err. (Ma)	Ft
Shell Canyon	BH11B	369	1497	22	0.64
	BH12	177	1673	11	0.74
	BH13B	103	1887	6	0.72
	BH13C	155	1887	9	0.75
	BH14	126	2027	8	0.72
	BH16	194	2405	12	0.79
	BH17	129	2716	8	0.76
Five Springs	BH5SPR	65	2012	4	0.76
	BH2	71	2448	4	0.74
	BH3	64	2603	4	0.53
South Gneiss	BH4B	65	2563	4	0.67
	BH8	62	2505	4	0.73
	BH10	80	2984	5	0.75
	BHM-739	77	2566	5	0.78
	BHM-508	68	2633	4	0.8
	BHM-508B	75	2633	4	0.78
	BHM-507	75	2929	5	0.78
East Side	BH18	72	1948	4	0.75
	BH18B	86	1948	5	0.78
	BH19B	89	2225	5	0.71
	BH19C	77	2225	5	0.77
	BH20B	88	2298	5	0.68
	BH21	81	2417	5	0.68
	BH22	112	2557	7	0.62
	BH30	126	2655	8	0.81
	BHM-725	95	2609	6	0.77
Central	BH23	108	3301	6	0.67
	BH23B	119	3301	7	0.64
	BH25	88	3139	5	0.52
	BH32B	100	3414	6	0.78
	BH33	86	2890	5	0.66
	BHM-618	70	2377	4	0.8

Table 1. (U-Th)/He ages of samples from all locations in the Bighorns. Samples with letters indicate replicate analyses.

RESULTS

(U-Th)/He ages for samples from all locations are shown in Table 1, and a plot of elevation vs. He age is given in Figure 2. Ages of samples from Shell Canyon range from as old as ~369 Ma to as young as ~126 Ma.

Samples taken from the lowest elevations at the western end of Shell Canyon yield the oldest ages, while those collected from the crest of the range at higher elevations give ages that generally cluster around ~120 Ma.

DISCUSSION

The Shell Canyon He ages exhibit an unusual age-elevation trend. According to the gently arched structural geometries proposed for the western side of the Bighorns near Shell Canyon (Figure 3), He ages of the Shell Canyon samples should be younger at lower elevations and gradually become older towards the top of the range. This pattern would be expected if the basement had been uniformly and deeply buried prior to the Laramide, and then during uplift simply bulged upwards beneath the overlying Paleozoic and Mesozoic sedimentary cover. In this scenario, exhumation of the uplifting block would have unroofed the central part of the range before the margins, and the higher samples (BH23, BH25, BH32) would have begun accumulating ^4He before the lower sample a relatively younger age. Ages calculated in this study, however, are oldest at the lowest elevations, (BH11, age 369 Ma) and younger towards the higher elevations at the top of the range (e. g. BH14 age 126Ma).

The He ages obtained in this study mandate that the lower, western side of the Bighorns must have occupied a high structural position since at least ~369 Ma, if not longer. The lower elevation sample, BH11, has been accumulating ^4He for at least ~369Ma, which indicates that the western slope of the Bighorns along Shell Creek has probably not experienced temperatures warmer than ~70°C since at least the early Paleozoic Era. In all likelihood, this region has probably been cooler than ~70°C since the Precambrian, as the calculated age of BH11 assumes a transition from complete ^4He loss to total ^4He retention at 369 Ma. In reality, this region of the basement may have resided in the ^4He Partial Retention Zone (PRZ) for much longer than the last 369 Ma (see Reiners and Crowley, this volume, for further explanation of the ^4He PRZ). Samples with younger He ages from the core of the range likely occupied deeper structural positions (see Figure 4, cross section of Shell Canyon, for details).

The presence of older He age samples at lower elevations than younger He age samples in Shell Canyon requires that the basement and the He PRZ in this region were never deeply buried, and were strongly folded during the Laramide. The Precambrian/Cambrian erosional surface across the Bighorns remains today as a north-south trending antiformal arch atop basement that folded as a result of Laramide crustal shortening, suggesting that the basement did flex ductilely as compressional deformation took place (see Figure 3 and Figure 4).

CONCLUSIONS

Apatite (U-Th)/He ages from Shell Canyon indicate that asymmetrical folding of the crystalline basement in this region occurred during the Laramide Orogeny. He ages also demonstrate that the far-western region of the basement in Shell Canyon has not experienced temperatures above ~70°C since at least 369 Ma due to extended




Figure 2. Plot of apatite (U-Th)/He ages vs. elevation above sea level.




Figure 3. Schematic cross-section through the Bighorns. after Stone, 1993.

shallow burial (no greater than 2 km) beneath pre-Laramide sediments. In addition, a colder than normal geotherm (~15°C/km) has probably existed through the crust in this region since the early Paleozoic. The higher elevation




Figure 4. Schematic cross-section through bottom of Shell Canyon, showing reconstructed paleo-PRZ at ~65 Ma. Thin black line denotes modern-day erosional surface. Apatite He ages are noted next to each sample. Elevation and horizontal distance units are given in meters. Vertical exaggeration is 3.

eastern end of Shell Canyon experienced burial beneath pre-Laramide sediments sufficiently thick enough to keep the basement heated to greater than ~70°C prior to about 120Ma. Variations in the thickness of sedimentary cover over the Bighorns likely produced the wide variability in measured apatite (U-Th)/He ages.

REFERENCES

- Crowley, P.D., Reiners, P. W. 2000. Isotopic Studies in the Bighorn Mountains, Wyoming: The U-Th/He Geochronology, Oxygen Isotope and Structural Perspectives. Thirteenth Keck Research Symposium in Geology Proceedings (Abstract Volume).
- Crowley, P.D., Reiners, P.W., Farley, K. A., Brown, H. E. Kaye, G.D., Reuter, J. M., 1999. Apatite (U-Th)/He Thermochronometry of the Bighorn Mountains, Wyoming: Implications for Structural and Sedimentation History. EOS, Transactions, American Geophysical Union Volume 80, Number 46, p. F1171.
- Darton, N.H., 1906, Geology of the Bighorn Mountains, USGS Professional Paper no. 51, 129pp.
- Farley, K.A., Wolf, R.A., and Silver, L.T., 1996, The Effects of Long Alpha Stopping Distances on (U-Th)/He Ages, *Geochimica et Cosmochimica Acta*, v. 60, p. 4223-4229.
- House, M.A., Wernicke, B.P., and Farley, K.A., 1998, Dating Topography of the Sierra Nevada, California, Using Apatite (U-Th)/He Ages, *Nature*, v. 396, p. 66-69.
- Reiners, P.W., Farley, K. A., Crowley, P.D., Brown, H. E. Kaye, G.D., Reuter, J. M., 1999. Apatite (U-Th)/He Thermochronometry of the Wind River and Bighorn Ranges: Contrasting Uplift and Exhumation Styles of Wyoming Laramide Uplifts. EOS, Transactions, American Geophysical Union Volume 80, Number 46, p. F 1035.
- Reuter, J.W., 2000. (U-Th)/He Thermochronometry in the Bighorn Mountains, Wyoming: Constraints on Timing of Exhumation. Thirteenth Keck Research Symposium in Geology Proceedings (Abstract Volume).
- Snoke, Arthur W., 1992. Geologic History of Wyoming Within the Tectonic Framework of the North American Cordillera. *Geology of Wyoming: State Geological Survey Memoir No. 5*, p. 2-56.
- Stone, D.S. 1993. Basement-Involved Thrust-Generated Folds as Seismically Imaged in the Subsurface of the Central Rocky Mountain Foreland. Schmidt, C. J., Chase, R. B., & Erslev, E. A. (editors), *Laramide Basement Deformation in the Rocky Mountain Foreland of the Western United States*, Special Paper - Geological Society of America, 280, p. 271-318.
- Wise and Obi, 1992, Laramide Basement Deformation in an Evolving Stress Field, Bighorn Mountain Front, Five Springs Area, Wyoming. *American Association of Petroleum Geologists Bulletin*, v. 76, No. 10, p. 1586-1600.
- Wolf, R.A., Farley, K.A., and Silver, L.T., 1996, Helium Diffusion and Low-Temperature Thermochronometry of Apatite, *Geochimica et Cosmochimica Acta*, v. 60, p. 4231-4240.

