

# SMALL SCALE FOLD AND FOLIATION ORIENTATIONS RELATED TO MULTIPLE STAGES OF DUCTILE DEFORMATION, QUETICO-WAWA BELT JUNCTION, NEAR REID LAKE, QUETICO PROVINCIAL PARK, ONTARIO, CANADA.

Bryan Shuman  
The Colorado College  
Colorado Springs, Colorado

Aaron Sheaffer  
Franklin and Marshall College  
Lancaster, Pennsylvania

## INTRODUCTION

The 1993 Quetico Keck project focused on understanding metamorphic terranes of the Canadian Shield, as they appear just north of the Minnesota-Ontario border within Quetico Provincial Park. The study area was located around Reid Lake. Sections of two subprovinces of the Superior Province of the Canadian Shield, the Quetico and the Wawa, and the junction between them, lie within the area. The Quetico subprovince has been interpreted as an accretionary wedge, consisting primarily of metagreywackes (Percival and Williams, 1989). It has undergone Amphibolite facies metamorphism which was overprinted by later Greenschist facies metamorphism (Oxboel, 1990; Percival, 1989). Related to this metamorphism, the Quetico subprovince has an axial zone of extreme anatexis migmatization and S-type plutonism (Card, 1990; Percival and Williams, 1989). The Wawa subprovince is a Granite-Greenstone belt, interpreted as a metamorphosed volcano-plutonic arc (Percival and Williams, 1989; Sylvester et al, 1987). Oceanic crust, north of the passive margin of the Wawa subprovince, was subducted beneath the Wabigoon volcano-plutonic arc. The Quetico subprovince was the fore-arc wedge of this system. When subduction ended, the Quetico collided with and sutured to the Wawa. This shifted the Quetico to a back-arc setting related to subduction on the southern margin of the Wawa (Williams, 1990; Card, 1990; Percival and Williams, 1989).

The rocks of the study area represent the suture boundary. Since these rocks were ductile and partially melted during the collision, the rocks of the two subprovinces are highly mixed together in the area. Thus no distinct boundary between the subprovinces exists. But in the zone of mixing, the suturing event created folds and foliations. This specific project was conducted to investigate the generations of these folds and foliations. Heavy anatexis of some units caused the study to be limited to the schist-dominated units of the area. In those units, thin leucosomes highlighted folding and schistosity was clear. The small size of the study area allows for the assumption that orientations, of axial surfaces and foliations, remained constant from outcrop to outcrop. Thus, orientations could be used to help distinguish between fabrics.

## FOLDS

The orientations of 85 axial surfaces and hinge lines were measured in the field. Variations in strike, dip, and plunge are not isolated to individual localities, but are widely distributed. Axial surface orientations differ within the same outcrop. This was due to the presence of three generations of folds in the Reid Lake area. Although these have different orientations, all were observed to have disharmonic, noncylindrical properties.

When the axial surfaces were plotted as poles on stereonet, the three distinct orientations appeared. The fold hinge lines, which correspond with these axial surface categories, were then plotted on the stereonet. A great circle was fit to the hinge lines to represent the local orientation of the axial surface. The poles to these great circles always fell within the tight cluster of measured axial surface poles. The orientation of the F<sub>1</sub> axial surface was N30E;60W. The F<sub>2</sub> great circle showed an axial surface orientation of N75E;75W. The F<sub>3</sub> axial surface orientation was N05W;55W.

Each of the three generations of folds has distinctive characteristics. F<sub>1</sub> was usually isoclinal folding of an S<sub>0</sub> compositional banding. The limbs of these folds are parallel with the F<sub>1</sub> axial surface, which runs subparallel to the Quetico-Wawa boundary. F<sub>1</sub> folds usually showed S-symmetry, although some Z and M folds were seen. F<sub>2</sub> was ptygmatic and limited isoclinal folding of the F<sub>1</sub> axial surface and S<sub>0</sub> around the F<sub>2</sub> axial surface. These F<sub>2</sub> folds showed only S-symmetry. F<sub>3</sub> appeared as ptygmatic and isoclinal folding of the limbs of F<sub>1</sub> and F<sub>2</sub> isoclinal folds around the F<sub>3</sub> axial surface. These folds also have S-symmetry. Often, though, F<sub>3</sub> folding appeared only as ductile elongation of the older foliations, such as the S<sub>0</sub> leucosomes. In horizontal outcrop surfaces, this appeared as normal separation of leucosome boundaries. The paradox of these folds is that F<sub>2</sub> ptygmatic folding appears on the limbs of F<sub>3</sub> folds. Some F<sub>4</sub> ptygmatic folds were also observed.

## FOLIATIONS

Up to six foliations have been observed in the outcrops of the Reid Lake area. Characteristics such as cross-cutting relationships and amount of post-formation deformation were used to assign numbers S<sub>0</sub> to S<sub>4</sub>. S<sub>0</sub>, as stated

would propagate from the southwest and the resulting duplex structure, once eroded, would account for the tonalite-amphibolite-tonalite sequence. In the SLF type section, brecciation dips more gently (25-35°NW) than the local S<sub>1</sub> foliation. This brecciated, shallowly dipping plane is suggestive of a thrust fault. However, applying present-day plate tectonics to Archean docking events may be problematic. The Archean continental crust was almost certainly substantially thinner, hotter, and therefore less dense than modern-day continental crust. Accretion and even obduction may not have taken place in the same manner as in the more recent geologic past. Convergence at subduction boundaries may not have been necessary. However, having found neither an isoclinal fold nose, nor evidence of an irregular tonalite-amphibolite contact, thrusting must be viewed as an acceptable hypothesis.

**CONCLUSIONS:** Despite the problems intrinsic in each of the three models, the repetition of tonalite at the south end of the SLF could be explained by any of them. The double layer model seems the most problematic. The thinness of the amphibolite and the conjectured regular contact do not accommodate the double layer model. The single layer model is more probable because of the isoclinal folding present within the amphibolite. These could be parasitic to larger regional isoclinal folds. No major fold noses were discovered within the study area, but this does not eliminate the possibility of the existence of one. Preliminary observations most strongly support a model based on thrust faulting. The shallowly-dipping brecciated zone could result from thrust faulting produced by compressional stresses. Regardless of which model is selected, field observations suggest that the SLL has undergone multiple deformational events; both ductile followed by brittle including the faulting along the Silence Lake Lineament.

### References

- Bauer, R.L. and Bidwell, M.E., 1990, Contrasts in the response to dextral transpression across the Quetico-Wawa subprovince boundary in northeastern Minnesota: *Canadian Journal of Earth Science*, V. 27, p. 1521-1535.
- Card, K.D. and Ciesielski, Andre, 1986, Subdivisions of the Superior Province of the Canadian Shield: *Geoscience Canada*, V. 13, p. 5-13.
- Percival, J.A. and Williams, H.R., 1989, Late Archean Quetico accretionary complex, Superior province, Canada: *Geology*, V. 17, p. 23-25.
- Southwick, D.L., 1991, On the genesis of Archean granite through two-stage melting of the Quetico accretionary prism at a transpressional plate boundary: *Geological Society of America Bulletin*, v. 103, p. 1385-1394.
- Woodard, H.H., 1992, Rock units and deformational structures related to the junction of the Quetico and Wawa subprovinces, Basswood Lake to Yum Yum Lake, Quetico Provincial Park, Ontario: Fifth Keck Research Symposium in Geology, Wooster Ohio, p. 70-75.
- Woodard, H.H., Root, S., Askren, D., 1994 Seventh Keck Research Symposium in Geology, Wooster Ohio.

above, is compositional banding. Usually these bands include leucosomes of plagioclase and quartz, in places with K-feldspar, biotite, and amphibole. There are also mesosomes of amphibole, with homogeneously incorporated feldspar, quartz, and biotite. Except in fold noses, isoclinal folding has transposed  $S_0$  banding into the  $F_1$  axial surface. Thus, the  $F_1$  axial surface is highlighted by this migmatite banding as  $S_1$ .  $S_1$  also appears as the primary schistosity in the mesosomes, as well as in the local tonalite sills.

The other four foliations appear similar to each other. Generally, they are observed as parallel alignment of individual crystals of amphibole, biotite, quartz, and plagioclase. They are not however, segregated bands of these minerals. These foliations also appear in intrusive dikes with graphic texture as parallel quartz grains. In other dikes and sills, including the tonalites, they occur as oriented amphiboles. However, in units which have experienced high levels of in situ melting, the occurrence of these foliations, and any folding, is very limited.

Three of these four foliations appear to run parallel to subparallel with the axial surfaces of folds.  $S_2$  runs about N70E, 80N - N70W, 80S, parallel with the  $F_2$  axial surface. The mineral alignments of  $S_2$  crosscut  $S_1$  in  $F_1$  folds. Measurements of  $S_2$  show that its orientation from place to place is more consistent than  $S_1$ . This suggests that  $S_1$ , more than  $S_2$ , has been rotated by later deformation.  $S_3$  runs about N15E, 70W - N15W, 70E, parallel with the  $F_3$  axial surface. It commonly cross-cuts  $S_2$  in  $F_2$  folds, and generally is shown to offset  $S_2$ . Also, the elongation of  $S_0/S_1$  leucosomes in the limbs of  $F_2$  folds is parallel with  $S_3$ . Measurements of  $S_3$  show that it has a more consistent orientation than  $S_1$  and  $S_2$ . This suggests that  $S_3$  has been rotated less than either  $S_1$  or  $S_2$ .  $S_4$  has been observed to offset  $S_3$ .  $S_4$  crosscuts  $F_2$  and  $F_3$  folds, running subparallel to  $S_1$  (N30E, 70W).  $S_1$  is folded by  $F_2$  and  $F_3$ . Therefore,  $S_4$  cannot be the same as  $S_1$ , but must be a younger foliation. Some  $F_1$  folds may actually be  $F_4$  folds since the  $F_1$  axial surface and  $S_4$  are parallel. The fourth of these foliations was only rarely observed preventing any determination of its age relationships.

## CONCLUSIONS

This information about the folds and foliations has allowed for a history of local deformation to be reconstructed. First, since  $S_0$  banding is mainly related to migmatites, migmatization must have occurred before  $D_1$ .  $D_1$  then followed producing  $F_1$  isoclinal folds and  $S_1$  foliation. Following this,  $D_2$  created  $F_2$  pygmatic and isoclinal folds, and  $S_2$  foliation. Next,  $D_3$  took place creating ductile extension of  $S_0/S_1$ . At this time,  $S_3$  foliation and  $F_3$  pygmatic and isoclinal folds were formed. Finally,  $D_4$  created at least  $S_4$  foliation and possibly  $F_4$  folds. Due to rotation, little can be said about the strain associated with the  $D_1$  and  $D_2$ . However, limited rotation of  $D_3$  structures suggests that its direction of greatest shortening, Z, was trending at N85E and plunging 35 degrees. The direction of greatest lengthening, X, was then in the plane of the axial surface trending at N05W and horizontal. However, it is not certain whether or not deformation was coaxial. The same is true for  $D_4$  deformation. There appears to have been no latter rotation of  $D_4$  structures. Thus for  $D_4$ , Z trended at S60E and plunged 20 degrees.

## REFERENCES

- Card, K. D., 1990, A review of the Superior Province of the Canadian Shield, a product of Archean accretion: *Precambrian Research*, v. 48, pp. 99-156.
- Oxboel, Lisa, 1990, Petrographic Study of Biotite Schist-rich Migmatite, Quetico Belt, Superior Province, Ontario, Canada: Third Keck Geological Research Symposium Abstracts Volume, pp. 115-118.
- Percival, John A., 1989, A regional perspective of the Quetico metasedimentary belt, Superior Province, Canada: *Canadian Journal of Earth Sciences*, v. 26, pp. 667-693.
- Percival, John A. and Williams, Howard R., 1989, Late Archean Quetico accretionary complex, Superior province, Canada: *Geology*, v. 17, pp. 23-25.
- Sylvester, Paul J. et al, 1987, Tectonic setting of late Archean bimodal volcanism in the Michipicoten (Wawa) greenstone belt, Ontario: *Canadian Journal of Earth Sciences*, v. 24, pp. 1120-1134.
- Williams, Howard R., 1990, Subprovince accretion tectonics in the south-central Superior Province: *Canadian Journal of Earth Sciences*, v. 27, pp. 570-581.




Figure 1. Poles of S1 (dots) and F1 fold axial surfaces (boxes)




Figure 2. Poles of S2 (dots) and F2 fold axial surfaces (boxes)




Figure 3. Poles of S3 (dots) and F3 fold axial surfaces (boxes)




Figure 4. A photograph-based sketch of a trondhjemite leucosome in amphibole-schist, on the west shore of Reid Lake, showing F2 folding. The fold limbs are offset by S3 and show F4 ptygmatic folding. In the schist, S3 mineral alignments are offset by S4. S2 mineral alignments run parallel to the F2 axial surfaces. Pen for scale.

## Author Index

Archuleta, LeAndra L.	60
Askren, Daniel R.	194
Barrera, John	177
Blair, J. Luke	101
Blakely, Christopher Todd	105
Brackin, Stephanie	199
Bradford, Andrew W.	109
Brady, John B.	54
Burger, H. Robert	9
Burger, H. Robert	54
Butler, Jessica	181
Cady, Pamela	64
Cheney, John T.	54
Coombs, Michelle L.	17
Curran, H. Allen	6
DeSimone, David	173
Donnelly, Kathleen E.	152
Elliott, Brent A.	199
Farthing, Dori	203
Fisher, Robin	67
Ford, Kirsten	206
Greene, Mary K.	156
Hanson, Lauren Polly	112
Holden, Jonathan B.	21
Howe, Evan S.	116
Humm, Amy	210
Jacob, Lisa	71
Johnson, Ann	119
Julia F. Daly	148
Ketcham, Brannon J.	123
King, Jonathan Tobias	75
Kozak, Samuel J.	13
Kroeger, Glenn C.	99
Leshner, Rebecca S.	127
Libbey, Laura	185
Lowell, Josh	78
Lubick, Naomi	25
Madera, Edwin	189
McCormick, Casey	206
Mendelson, Carl V.	6
Nashem, William D.	131
Newton, Robert M.	173
Parkins, Rebecca Kern	135
Peck, William H.	82
Petcovic, Heather	181
Pierson, Ana	185
Plummer, Benjamin P.	29
Poulsen, Chris	86
Root, Samuel I.	99
Root, Samuel I.	194
Rosen, Joy	177
Russell, Elizabeth M.	160

Saffer, Demian	189
Sak, Peter	210
Savina, Mary E.	4
Savina, Mary E.	144
Schuh, Kathleen J.	33
Seckler, Michael S.	37
Sheaffer, Aaron	214
Shuman, Bryan	214
Sincock, Mary Jennifer	90
Spencer, Edgar W.	99
Stewart, Brooke	203
Sturm, Marnie	41
Symchych, Elizabeth M.	45
Tierney, Kara A.	94
Titzel, C. Scott	139
Vanden Bergh, Frederick J.	49
Wiebe, Robert A.	13
Williams, Julianne M.	164
Wilson, Robert	168
Wobus, Reinhard A.	13
Woodard, Henry H.	1
Woodard, Henry H.	194

